

DESCRIPTION OF SURFICIAL-GEOLOGIC MAP UNITS

(Map units below might not all appear on this sheet)

This map shows the distribution of unconsolidated deposits and undifferentiated bedrock exposed at the surface in part of the central segment of the proposed natural-gas pipeline corridor straddling the Alaska Highway from Robertson River to Tetlin Junction in the Tanacross Quadrangle. Units were mapped by interpretation of false-color infrared ~1:65,000-scale aerial photographs taken in July 1978, August 1980, and July 1983 and verified by field checking in 2007 and 2008.

Map units shown in parentheses such as (Qc1), indicate combination map units consisting of bedrock overlain by thin to discontinuous material of the map unit shown.

UNCONSOLIDATED DEPOSITS

- UNDIFFERENTIATED FLOODPLAIN ALLUVIUM**—Chiefly well sorted and well stratified polygenetic pebble gravel, sand, and silt comprising channel and overbank deposits of generally small streams; unfrozen to discontinuously frozen with low to moderate ice content
- ACTIVE FLOODPLAIN ALLUVIUM**—Chiefly well sorted and well stratified polygenetic pebble gravel, sand, and silt with rare to scattered cobbles comprising river bars subject to recurrent inundation by streams every 5 yrs or less (Chapin and others, 2006); mapped extent is a function of river level (stage) and reflects the transitory extent of exposed river bars at the time the photographs were taken; in braided and anastomosing reaches, active channels typically shift positions from year to year and present channel locations may differ from locations in the photography on which the deposits were mapped; active alluvium underlies upper stream bank and active stream channels and includes point-bar and meander-scut deposits (Brakenridge, 1988); composed dominantly of gravel and sand where stream is braided and anastomosing and sand and silt bars and cover deposits where meandering; prone to liquefaction where fine grained and unfrozen (Harp and others, 2005); where braided, subject to formation of extensive, thick seasonal-stream incises (auflages); generally unfrozen, except seasonally frozen to depth of frost penetration; shallow water table
- ABANDONED FLOODPLAIN ALLUVIUM**—Chiefly 10 to 20 ft (3 to 6 m) of overbank sandy silt and siltly sand overlying sandy, polygenetic riverbed gravel beneath surface with widespread low and level sand dunes and subject to stream flooding about once every 500 to 1,000 yrs (Mann and others, 1995); may include several surfaces at different levels; overbank sequences include flood-related features, like natural levees, crevasse-splays, and expansion fans near channels and fine-grained pebbly swale deposits farther from channels (Brakenridge, 1988; Mann and others, 1995); may contain organic-silt channels (1 to 7 to 20 ft (0.3 to 6 m) thick; surface pebbles generally discontinuous and widespread in backwater areas away from channels; floodplain lakes are larger than lakes on younger floodplains and typically have rounded to scalloped shorelines formed by thermokarst erosion; generally frozen with low to moderate ice content
- ALLUVIAL-FAN DEPOSITS**—Fan-shaped deposits of unsorted to well sorted gravel, sand, and silt with numerous cobbles and boulders in proximal zone; lithologies reflect bedrock of source area; in general, size of clasts decreases and degree of sorting increases downslope; typically mixed with debris-flow deposits in proximal part of fans; unfrozen to discontinuously frozen, except in fine-grained distal zones where permafrost may be shallow and continuous; ice content low to moderate
- INACTIVE FLOODPLAIN ALLUVIUM**—Chiefly 2 to 20 ft (0.6 to 6 m) of overbank siltly sand and sandy silt overlying gravelly, polygenetic riverbed sand and gravel beneath surface subject to flooding as often as twice to ten times per century (Mann and Reger, 1991; Yarie and others, 1998; Chapin and others, 2006); may include more than one surface at different levels; overbank sequences include flood-related features such as natural levees, crevasse-splays, and expansion fans near channels, and fine-grained back-levee swale deposits farther from channels (Brakenridge, 1988; Mann and others, 1995); small lakes have linear, arcuate, and confluent outlines (Webster and Pevé, 1961, 1970; Pevé, 1970; Reger and Hubbard, 2009); surface pebbles generally absent; prone to liquefaction where fine grained and unfrozen (Harp and others, 2005); generally unfrozen in younger areas and discontinuously frozen in older areas with low to moderate ice content; active channels may be underlain by 5 to 20 ft (1.5 to 6 m) of generally unfrozen sand and silty sand; of inactive channels may include 7 to 12 ft (2.1 to 3.6 m) of discontinuously frozen sand and silt with moderate to high ice content over sand and gravelly sand
- STREAM-TERRACE ALLUVIUM**—Chiefly 4 to >20 ft (>1.2 to >6 m) of organic sandy silt and siltly sand overlying well sorted, polygenetic sand and gravel beneath stream terrace troughs no longer subject to inundation by the stream that deposited the alluvium (Kreig and Reger, 1982); may include several levels and flood-related features, like natural levees, crevasse-splays, and expansion fans near channels; may incorporate overwash alluvium of Donnelly age in highest terraces; locally covered by <15 ft (<4.5 m) of lowland loess and collan-sand blanket and dune complexes, especially close to active sediment sources; trace lakes with rounded to scalloped shorelines formed by thermokarst erosion are typically present (Webster and Pevé, 1961, 1970; Pevé, 1970; Reger and Hubbard, 2009); locally subject to seasonal stream incises where building of auflages in stream channels diverts subsequent drainage and spreads arifacts and meltwater across terraces that would not otherwise be flooded (Springer and others, 1976; Sloan and others, 1976); continuously to discontinuously frozen with low to moderate ice content
- FLOOD DEPOSITS**—Eruption fans, crevasse-splay deposits, pendant bars, and linear bars fining away from the modern floodplain of the Tanana River or its tributaries; generally unfrozen to discontinuously frozen with low to moderate ice content
- SLACKWATER FLOOD DEPOSITS**—Chiefly organic sandy and siltly backwash sediments deposited during floods in slackwater basins separated from main stream by expansion fans and natural levees and crevasse-splays; unfrozen to discontinuously frozen with low to moderate ice content
- TERRACE DEPOSITS OF YOUNGER TOK FAN**—Surface above inactive and abandoned floodplains of Tok River displays former meandering and anastomosing drainage channels of Tok River; composed of micaceous cover silt with trace clay up to 5 to 12.7 cm thick overlying poorly sorted, generally massive to crossbedded, matrix-supported pebbly medium-to-course sand with trace silt and rare polygenetic cobbles up to 4 in (10.2 cm) diameter; moderate imbrication; depth of carbonate-bottomed pebbles varies up to 32 in (0.8 m); carbonate cementa and coarse sand to bottom of pebbles; silt cap discontinuous and 0.1 to 0.25 cm thick; matrix color dark yellowish brown (10YR4/6) to grayish brown (10YR5/2); locally poorly drained; discontinuously frozen with low to moderate ice content
- ZONE OF GROUNDWATER EMERGENCE ON OLDER TOK FAN**—Surface features on typically well drained western, older Tok fan that indicate emergence of groundwater include swampy vegetation, peat, and standing surface water; the presence of water in shallow, alluvial trenches; networks of shallow drainage channels originating at drainage springs; and a concentration of clearwater ponds and lakes, and thermokarst pits; inferred to be continuously frozen and ice-rich
- SLACKWATER FLOOD DEPOSITS**—Chiefly organic sandy and siltly backwash sediments deposited during floods in slackwater basins separated from main stream by expansion fans and natural levees and crevasse-splays; unfrozen to discontinuously frozen with low to moderate ice content
- TERRACE DEPOSITS OF YOUNGER TOK FAN**—Surface above inactive and abandoned floodplains of Tok River displays former meandering and anastomosing drainage channels of Tok River; composed of micaceous cover silt with trace clay up to 5 to 12.7 cm thick overlying poorly sorted, generally massive to crossbedded, matrix-supported pebbly medium-to-course sand with trace silt and rare polygenetic cobbles up to 4 in (10.2 cm) diameter; moderate imbrication; depth of carbonate-bottomed pebbles varies up to 32 in (0.8 m); carbonate cementa and coarse sand to bottom of pebbles; silt cap discontinuous and 0.1 to 0.25 cm thick; matrix color dark yellowish brown (10YR4/6) to grayish brown (10YR5/2); locally poorly drained; discontinuously frozen with low to moderate ice content
- ZONE OF GROUNDWATER EMERGENCE ON YOUNGER TOK FAN**—Surface features on the eastern, younger Tok fan that indicate emergence of groundwater include swampy vegetation, peat, standing surface water, and networks of shallow drainage channels

COLLUVIAL DEPOSITS

- UNDIFFERENTIATED COLLUVIUM**—Blankets, aprons, cones, and fans of heterogeneously mixed angular to subangular rock fragments, gravel, sand, and silt formed by creep, gravity-driven mass movements involving sliding, flowing, gelifraction, and frost creep of weathered bedrock and modified glacial drift; cobbles and boulders are scattered to numerous; lower headwalls of cirques and upper walls of glaciated valleys include talus aprons, incipient rock glaciers, and related features; as well as steep fine-grained talus by snow avalanches and debris flows; may include thin residual deposits and lags of former Tertiary bedrock and highly modified drift of earlier glaciations; low-gradient, western, older part of broad Tok fan, which is composed of clay- and matrix-supported, tabular, massive to crudely bedded gravels interbedded with minor beds of crudely bedded pebbly sand; beds average ~3.5 ft (1 m) thick, parallel the fan surface, and contain rare extraordinarily large flood boulders; unfrozen to discontinuously frozen; low ice content
- SNOW-AVALANCHE DEPOSIT**—Steep fans of heterogeneous pebbly debris with some gravel, sand, and silt deposited by snow avalanches in and downslope of cirques in steep alpine settings; angular rock debris with scattered, small clasts of granitic rock; debris flow debris with the largest fragments farther downslope; typically associated with talus cones and aprons; discontinuously frozen with low to moderate ice content
- DEBRIS-FLOW DEPOSIT**—Chiefly tongues of angular rock fragments and coarse gravel with a sandy matrix deposited on steep colluvial slopes and fans and in rock-walled upper stream valleys by flowing slurries of mud, sand, rock debris, and gravel generated during sudden intense rainstorms; initial fracturing and later winnowing, leaving coarse gravel and rubble tongues and lobes some with natural talus of cobbles and boulders up to 7 ft (2.1 m) high bounding medial channels with relict U-shaped cross profiles measuring 70 to 80 ft (3 to 21 m) across and 10 to 60 ft (3 to 18.3 m) deep; many large boulders and blocks have small debris mounds and scattered cobbles on upper surfaces; generally unfrozen to discontinuously frozen with low ice content
- MIXED COLLUVIUM AND ALLUVIUM**—Primarily fan-shaped or oblong, massive to poorly stratified, generally inorganic silt mixed with sandy angular to subangular pebbly gravel derived from weathered bedrock uplands and loess-covered moraines, and laid down by debris flows and hyperconcentrated flows produced during brief, intense local summer storms; colluvial processes > fluvial processes; surface slightly irregular; contains numerous cobbles in glacial terraces and angular to subangular, fresh to weathered rock fragments and gran in weathered granitic bedrock terrain; discontinuously to continuously frozen with low to moderate ice content
- TECTONICALLY DEFORMED COLLUVIAL-FLUVIAL DEPOSITS**—Arcuate ridges of poorly stratified, coarse, sandy gran fragments with trace silt, numerous pebbles and scattered subrounded to rounded granitic boulders up to 9.2 ft (2.8 m) diameter initially deposited as piedmont aprons southwest of Tanacross (Airtfield sheet 3) by debris flow derived from the steep mountain valley to the southwest and later tectonically deformed (Carver and others, 2010); sandy granitic matrix, color dark brown (5.5YR4/4) to light olive brown (2.5Y5/4); surface smoothly rounded with slopes between ~0° and ~17°; partially eluvial granitic boulders stand up to ~5 ft (1.5 m) in relief; heights of surface boulders greater where surface slopes are steeper; surface stepped by ~20° to ~25° sharp of shallow, local slope failures; discontinuously frozen with low ice content
- ROCK-GLACIER DEPOSITS**—Tongue-shaped heterogeneous surface blanket of angular to subangular blocks of local bedrock overlying deformed ice with trace to some gravel, sand, and silt at depth; where active, blocky surface layer is disrupted on steep marginal slopes and core debris is exposed; accumulated on floors and lower walls of cirques and glaciated valleys by flow of rock glaciers derived from shrinking of former glaciers (ice core) or from deposition, cementation, and deformation of precipitation-derived ground ice (ice cementation); surface typically has furrows, nested arcuate ridges arranged convexly downslope, and pits, and may have prominent lateral ridges; perennially frozen where active with moderate to high ice content
- LANDSLIDE DEPOSITS**—Lunate to triangular or fan-shaped, heterogeneous mixtures of large fractured bedrock blocks and pebble gravel with scattered to numerous cobbles and boulders and trace to some sand and silt deposited by near-surface to deep creeping, flowing, and sliding of failed bedrock and unconsolidated surficial deposits; surface features include gaging ground cracks where active, slight irregularities, hummocks, low longitudinal ridges, and terminal ledges; unfrozen to continuously frozen with low to moderate ice content
- ROCK-FALL DEPOSITS**—Rubble blanket or apron of large, angular rock fragments of local bedrock formed by collapse of upslope outcrop; unfrozen to discontinuously frozen with low ice content
- TALUS**—Cone- and apron-shaped heterogeneous mixtures of frost-riveted, angular rock fragments downslope of bedrock outcrops with trace to some gravel, sand, and silt deposited on steep bedrock slopes and at the mouths of steep bedrock outcrops by U-shaped profiles by snow avalanches, free fall, tumbling, rolling, and sliding; surface steep, slightly irregular, and covered with numerous rock fragments, particularly in distal zones; includes debris-flow tongues; blocks and boulders covered by crustose lichens where stable and lichens free where freshly displaced; unfrozen to discontinuously frozen with low ice content

EOLIAN DEPOSITS

- UNDIFFERENTIATED EOLIAN DEPOSITS**—Chiefly well sorted, massive to finely bedded, primarily airfall eolian sand and loess forming a blanket over bedrock ridges and hills and lowlands in the southern Yukon-Tanana Upland; complex stratigraphy may include retransported sand and silt; discontinuously to continuously frozen with low to high ice content
- LOESS**—Silt with up to 15 percent very fine sand carried by winds and deposited as a blanket over downwind topography (Pevé, 1981, 1985); mixed with eolian sand on lower slopes close to floodplain sources; may include intimate mixtures with retransported silt; thickness ranges from >20 ft (>6 m) close to active sediment sources to ~2 ft (<0.6 m) elsewhere (Lindholm and others, 1959); typically filled where >3 ft (>0.9 m) thick on steep upper slopes, but areas of mapped loess should be considered minimal because fills are locally obscured by dense vegetation cover; organic rich on lower slopes and lowland sites; moderate to high moisture content (~15 percent) in lowlands (Kreig and Reger, 1982); generally unfrozen, except discontinuously frozen with moderate to high ice content on some lower, south-facing slopes and continuously frozen and ice rich on some lower north-facing slopes and lowland sites
- RETRANSPORTED SILT AND SAND COMPLEXLY MIXED WITH LOWLAND LOESS**—Chiefly massive to well stratified organic silt and sandy silt with lenses and tongues of locally derived gravel and scattered to numerous angular rock fragments (particularly in upper valleys of small ephemeral streams) in loess areas and organic fine sand and silt deposited by near-surface to deep creeping, flowing, and sliding of failed bedrock and unconsolidated surficial deposits; surface features include gaging ground cracks where active, slight irregularities, hummocks, low longitudinal ridges, and terminal ledges; unfrozen to continuously frozen with low to moderate ice content
- EOLIAN SAND**—Chiefly blankets and dunes of fine to medium, massive to cross-bedded eolian sand with trace to some silt (Kreig and Reger, 1982, p. 9); dunes stand 5 to 15 ft (1.5 to 4.5 m) in relief and ridges may extend for up to 3 mi (4.8 km) in the direction of dominant summer winds; mapped extent, based on the presence of dunes, should be considered minimum; dune ridges locally crown steep slopes that are the sand source; discontinuous with thicknesses up to ~25 ft (~7.6 m); unweathered color dark brown (2.5Y5/2); generally covered by 1 to 3 ft (0.3 to 0.9 m) of loess (Lindholm and others, 1959); locally being deposited along the margins of braided floodplains; average moisture content ~8 percent (Kreig and Reger, 1982); discontinuously frozen with low to moderate ice content




GLACIAL DEPOSITS

- TILL AND ASSOCIATED MORAINAL DEPOSITS OF POST-DONNELLY GLACIATION**—Heterogeneous, non-stratified, polygenetic pebble-cobble gravel with some sand and silt and numerous subangular to subrounded boulders deposited by glacial ice and associated colluvial processes in upper mountain valleys during Holocene time; boulders of younger deposits are unvegetated or bear crustose lichens; older moraines are typically covered with tundra; loess cover thin and patchy to nonexistent; ice cores may be present, especially in younger moraines; unfrozen to discontinuously frozen with low to moderate ice content

MAP SYMBOLS

(Map symbols might not all appear on this sheet)

- PHOTONTERPRETTED CONTACT—Dashed where approximately located
- QUESTIONABLE IDENTIFICATION
- ACTIVE HIGH ANGLE FAULT—Dashed where approximately located, dotted where concealed
- Arrows indicate apparent direction of relative movement
- U, yellowish block; D, downwash block (Carver and others, 2010)
- ACTIVE THRUST FAULT—Dashed where approximately located, dotted where concealed
- hatched on upper plate (Carver and others, 2010)
- ANTIFORM—Dashed where approximately located, dotted where concealed (Carver and others, 2010)
- SP-1 ● LOCATION OF RADIOCARBON SAMPLE DISCUSSED IN TEXT
- SP-1 ● LOCATION OF SOIL PIT DISCUSSED IN TEXT
- V-1 ● LOCATION OF VEINTRACE SITE DISCUSSED IN TEXT
- A ● MAP LOCALITY DISCUSSED IN TEXT
- A—A' ● GEOLOGIC PROFILE DISCUSSED IN TEXT



SURFICIAL-GEOLOGIC MAP, ALASKA HIGHWAY CORRIDOR, PARTS OF THE TANACROSS A-5 and B-5 QUADRANGLES, ALASKA

by
R.D. Reger¹, T.D. Hubbard² and G.A. Carver³

2011

¹ Reger & Geologic Consulting, PO Box 3126, Soldotna, Alaska 99669-3126
² Alaska Division of Geological & Geophysical Surveys, 3354 College Road, Fairbanks, Alaska 99709-3707
³ Carver Geologic Inc., PO Box 52, Kodiak, Alaska 99615-0052

- TILL AND ASSOCIATED MORAINAL DEPOSITS OF DONNELLY GLACIATION**—Heterogeneous, nonstratified, polygenetic pebble-cobble gravel with some sand and silt and few to numerous subangular to subrounded boulders deposited by glacial ice and locally reworked by meltwater washing and associated mass-movement processes; may include esker and kame deposits; moraine relief 50 to 175 ft (15.2 to 53.3 m); kettle frequency ~16/m² (~6.4/km²); kettle fillings of silt, peat, and siltly colluvium generally thin but may be several feet (meters) thick close to active sources of eolian deposits; maximum till thickness ~500 ft (152.4 m); surface weathering profiles 1.5 to 2.5 ft (0.45 to 0.8 m) thick; friable sand matrix weathered to brown (10YR5/3); 25 to 35 percent of schists clasts are intact in weathering profiles and granitic clasts are fresh to slightly weathered; silt caps generally ~1 mm thick; discontinuous cover of loess generally ~3 ft (0.9 m) thick and weathered yellowish brown (10YR5/8) to light yellowish brown (10YR6/4) but eolian sand and silt mantle may be ~20 ft (6 m) thick close to active sediment sources and may obscure primary surface morphology; ventifacts exhibit slight to moderate surface polish and shallow pitting but lack facets and keels in lags developed beneath loess covers; ice-wedge casts generally rare and up to 3 ft (0.9 m) wide; unfrozen to discontinuously frozen with low to moderate ice content (Pevé and Holmes, 1964; Holmes, 1965; Carter and Galloway, 1978; Pevé and Reger, 1983a, table 3)
- TILL AND ASSOCIATED MORAINAL DEPOSITS OF DELTA GLACIATION**—Heterogeneous, nonstratified, polygenetic pebble-cobble gravel with some sand and silt and few to numerous subangular to subrounded boulders deposited by glacial ice and massive, sandy pebble gravel with rare cobbles deposited by glacial meltwater and associated mass-movement processes; may include esker and kame complexes; moraine relief 25 to 225 ft (7.6 to 68.6 m); kettle frequency ~3/m² (~1.2/km²); kettle fillings of silt, peat, and siltly colluvium may be several feet (meters) thick; maximum till thickness ~200 ft (~60 m); surface weathering profiles generally 3-7 ft (0.9-2.1 m) deep, on high-level surfaces may locally be ~10 ft (3 m) deep; friable to strongly cemented with numerous clasts; sand matrix weathered light yellowish brown (10YR6/4) to brownish yellow (10YR6/6); 1 to 10 percent of schist clasts are intact in weathered profiles and ~50 percent of granitic clasts are partially decomposed; silt caps range from ~0.04 to 0.12 in (~1 to ~3 mm) thick; discontinuously mantled by thin eolian sand and loess; loess cover weathered to light reddish brown (5YR6/4) (rubification); well-formed faceted and keeled ventifacts common in surface lags beneath loess covers; ice-wedge casts scattered to numerous and up to ~5 ft (1.5 m) wide; wedge fillings include deformed eolian sand that is locally pebbly; unfrozen to discontinuously frozen with low to moderate ice content (Pevé and Holmes, 1964; Holmes, 1965; Carter and Galloway, 1978; Pevé and Reger, 1983a, table 3)
- UNDIFFERENTIATED GLACIAL DRIFT OF PRE-DELTA GLACIATION**—Thin, discontinuous to continuous sheets of heterogeneous pebble gravel, sand, and silt with rare to numerous cobbles, boulders, and blocks up to 8 ft (2.4 m) in diameter deposited directly from melting glaciers and reworked by meltwater streams; includes drift of Durling Creek age and perhaps other older glaciations on upper to middle surfaces and lower mountain slopes south of Tanana River; sandy matrix weathered pale brown (10YR6/3) to brown (10YR5/3); surface morphology extensively modified by mass-movement processes; unfrozen to discontinuously frozen with low to moderate ice content (Pevé and Reger, 1983a; Weber, 1986; Dak-Rodkin and others, 2004)

GLACIOFLUVIAL DEPOSITS

- OUTWASH OF POST-DONNELLY GLACIATION**—Massive to well sorted, polygenetic pebble-cobble gravel with some sand and numerous subrounded to angular boulders deposited by meltwater streams from Holocene glaciers in upper mountain valleys; locally includes deposits of debris flows and rockfalls; clasts are generally fresh; surfaces unvegetated to vegetated with thin tundra; loess cover nonexistent to thin and patchy; unfrozen to discontinuously frozen with low ice content
- OUTWASH OF LATE DONNELLY AGE**—Coarse outwash gravel in steep-walled, flat-bottomed, broad channel incised into surface of outwash fan of Robertson River glacier north of Tan Lake; connects with kame-esker deposits in the southeastern corner of Corridor Segment 1 (Reger and others, 2008, sheet 2)
- OUTWASH OF DONNELLY GLACIATION**—Massive to well sorted, polygenetic pebble gravel with some sand and scattered to numerous subrounded to subangular cobbles and boulders ~7 ft (2.1 m) in diameter in proximal zones; surface weathering profiles ~3 ft (0.9 m) deep; sand matrix color varies from pale brown (10YR6/3) to brown (10YR5/3); 5 to 10 percent of foliated siltstones are typically split by frost action and granitic thinstones are fresh to slightly weathered in weathered profiles, except locally, where foliated thinstones are shattered to small, platy fragments and granitic clasts are reduced to crumbly remnants by the growth of calcite (calcite) in the upper 3 to 4 ft (0.9 to 1.2 m) of the outwash deposit; silt caps thin and discontinuous; cover sands discontinuous and up to ~10 ft (~3 m) thick; average loess cover ~0.4 ft (~0.1 m) thick and generally weathered light yellowish brown (10YR6/4) to brown (10YR5/3), except red (2.5YR5/6) where strongly oxidized after repeated wildfires (Pevé and others, 2006); ventifacts exhibit slight to moderate surface polish and pitting but no facets or keels in lags developed beneath loess covers; ice-wedge casts generally rare, but locally common and ~3 ft (0.9 m) wide (Pevé and Reger, 1983a, p. 62-66); deformed wedge fillings composed of brown to greenish gray silt with trace to some pebble gravel and scattered cobbles; unfrozen to discontinuously frozen with low ice content
- OUTWASH OF DELTA GLACIATION**—Massive to well sorted, polygenetic pebble gravel with some sand and numerous subrounded to subangular cobbles and boulders ~3.5 ft (~1.1 m) in diameter; coarse in proximal zones and fine where distal; surface weathering profiles ~2 ft (0.6 m) deep; sand matrix color varies from pale brown (10YR6/3) to very pale brown (10YR7/4); ~50 percent of foliated and granitic clasts in weathered profile are rotten; silt caps on clasts in weathered profile ~0.04 to 0.12 in (~1 to ~3 mm) thick; loess cover sands discontinuous and up to ~10 ft (3 m) thick; loess deep typically 1 to 2 ft (0.3 to 0.6 m) thick; well formed faceted and keeled ventifacts common in surface lags beneath loess covers; quartz pebbles in lags stained yellowish brown (10YR5/4) to very pale brown (10YR6/4); ice-wedge casts scattered to numerous and ~3.5 ft (~1.1 m) wide; deformed wedge fillings are typically eolian sand with trace to some silt and pebble gravel and may include scattered pebble ventifacts; unfrozen to discontinuously frozen with low ice content

LACUSTRINE DEPOSITS

- LAKE-BOTTOM DEPOSITS**—Chiefly silt and clay with some sand and organic material deposited in ephemeral lakes in backwater areas of inactive floodplains and behind ice-observed remnants in large lakes; discontinuously to continuously frozen with moderate to high ice content
- DELTA DEPOSITS**—Chiefly sand and silt with some organic material deposited in a lake basin by a stream entering the lake; during floods of the Tanana River, streams normally draining the lake into the river reverse direction and carry floodwaters and sediments into the lake basin; sporadically frozen with moderate to high ice content
- DEPOSITS OF ICE-SHOVED RIDGES**—Single or multiple 3 to 5-ft-high (0.9 to 1.5-m-high) ridges parallel to and 2 to 15 ft (0.6 to 4.5 m) above modern lake shorelines; composed of overwashed and severely and complexly deformed deposits of adjacent lake bottoms, including fine to coarse elastic lake-bottom sediments and peat with thin interlayered light gray lacustrine sands; built by shoreward transport of lake-bottom sediments by wind-driven, drifting lake ice (Pevé and Reger, 1983a, figs. 22A and B); unfrozen to discontinuously frozen with low to moderate ice content

PALUDAL DEPOSITS

- SWAMP DEPOSITS**—Primarily fibrous and locally woody, anoxic peat with organic silt and sand deposited in lowland sites (Kreig and Reger, 1982); ~8 ft (~2.4 m) thick; discontinuously to continuously frozen with moderate to high ice content

RESIDUAL DEPOSITS

- BLOCK RUBBLE**—Nests and blankets of angular to subangular blocks derived from frost wedging and jacking of underlying bedrock (autochthonous block fields) on high-level surfaces (Heimann of Carter, 2004) and/or as lags left by winnowing of sandy matrix from gelifraction deposits or thin till by subterranean piping (allochthonous block fields); locally may be included in units of thinly covered bedrock and in shallow strath terraces; sizes of blocks are function of joint spacing in local bedrock; associated microficial features formed by frost action and mass movement include stone polygons, stone nets and circles, stone stripes, nested circles and hummocks, and soil lobes and benches; frost jacking locally active; discontinuously frozen with low to moderate ice content

BEDROCK

- UNDIFFERENTIATED BEDROCK**—Outcrops of igneous, metamorphic, and sedimentary rocks; linear and curvilinear shallow troughs and linear changes of surface vegetation indicate the presence of planar bedrock structures
- THINLY COVERED BEDROCK**—Subcrops with <3 ft (<0.9 m) of loess cover; bedrock structures recognizable through thin veneer of surficial debris
- Complex map unit consisting of bedrock outcrops and thinly buried subcrops that cannot be mapped separately

¹ Estimated contents of sand and silt, based on field observations, are indicated by the terms "trace" and "some." "Trace" implies a general composition of 4 to 10 percent. "Some" implies a general composition of 10 to 30 percent. Estimated compositions of present are not recorded in the field. Terms used to describe the estimated percentages of cobbles and boulders are "numerous," "scattered," and "rare." "Numerous" implies that drilling through the deposit would encounter two cobbles or boulders in an interval of 2 ft (0.6 m); "scattered" implies that drilling would encounter two cobbles or boulders in an interval of 10 ft (3 m) to 15 to 45 m; "rare" implies that drilling would encounter two cobbles or boulders in an interval of 15 ft (4.5 m).

REFERENCES

Brakenridge, G.R., 1988, River flood regime and floodplain stratigraphy, in Baker, V.R., Kochel, R.C., and Patton, P.C., eds., Flood geomorphology: New York, John Wiley & Sons, p. 129-142.

Carter, P.E., 2004, Surficial geologic map of the Tanacross B-5 Quadrangle, east-central Alaska: U.S. Geological Survey Scientific Investigations Map 2850, version 1.0, 9 p., 1 sheet, scale 1:63,360.

2004, Surficial geologic map of the Tanacross B-5 Quadrangle, east-central Alaska: U.S. Geological Survey Scientific Investigations Map 2856, version 1.0, 9 p., 1 sheet, scale 1:63,360.

Carter, J.D., and Galloway, J.P., 1978, Preliminary engineering geology maps of the proposed natural gas pipeline route in the Tanana River valley, Alaska: U.S. Geological Survey Open-File Report 78-79, 29 p., 2 sheets, scale 1:125,000.

Carver, G.A., Benin, S.P., Solt, D.N., Castagnone, S.R., and Oshemiller, K.E., 2010, Active and potentially active faults in or near the Alaska Highway corridor, Robertson River to Tetlin Junction, Alaska Division of Geological & Geophysical Surveys Preliminary Interpretive Report 2010-1, 42 p.

Chapin, F.S., III, Viereck, L.A., Adams, P.C., Van Cleve, Keith, Fostie, C.L., Ott, R.A., Mann, Daniel, and Johnson, J.F., 2006, Successional processes in the Alaskan boreal forest, in Chapin, F.S., III, Osowood, M.W., Van Cleve, Keith, L.A., and Verbyla, D.L., eds., Alaska's changing boreal forest: New York, Oxford University Press, p. 100-120.

Coom, J.E., 1986, Rheologic, geomorphic, and sedimentologic differentiation of water floods, hyperconcentrated flows, and debris flows, in Baker, V.R., Kochel, R.C., and Patton, P.C., eds., Flood geomorphology: New York, John Wiley & Sons, p. 113-122.

Dak-Rodkin, Alejandra, Barnedreg, R.W., Friesse, D.G., Weber, Frances, Enkin, Randy, Smith, R., Zarka, G.D., Waters, Pamela, and Klassen, Rody, 2004, Timing and extent of Pleistocene glaciations in northwestern Canada and east-central Alaska, in Ehlers, J., and Gibbard, P.L., eds., Quaternary glaciations—extent and chronology, part II: North America, Van Nostrand Reinhold, New York, Elsevier, Developments in Quaternary Sciences, v. 2, p. 313-345.

Harp, E.L., Jibson, R.W., Kyeen, R.E., Keefer, D.K., Sherrill, B.L., Carver, G.A., Collins, B.D., Moss, R.E.S., and Sitar, N., 2003, Landslides and liquefaction triggered by the 11.75 Denali Fault earthquake of 2 November 2002: *Geology*, v. 31, no. 8, p. 1144-1149.

Holmes, G.W., 1965, Geologic reconnaissance along the Alaska Highway, Delta River to Tok Junction, Alaska: U.S. Geological Survey Bulletin 1181-H, 19 p., scale 1:125,000, 1 sheet.

Kreig, R.A., and Reger, R.D., 1982, Air-photo analysis and summary of landform soil properties along the route of the Trans-Alaska Pipeline System: Alaska Division of Geological & Geophysical Surveys Geologic Map 66, 149 p.

Lindholm, G.F., Thomas, L.A., Davidson, D.T., Handy, R.L., and Roy, C.J., 1959, Silt near Big Delta and Fairbanks, in Davidson, D.T., and Roy, C.J., eds., The geology and engineering characteristics of some Alaskan silts: U.S. Geological Survey Engineering Experiment Station Bulletin 186, p. 33-70.

Mann, D.H., Fostie, C.L., Rowland, E.L., and Bigelow, N.H., 1995, Spine succession, disturbance, and geomorphology on the Tanana River floodplain, Alaska: *Ecosystems*, v. 2, no. 2, p. 184-199.

Mason, O.K., and Bepko, J.E., 1991, Late Holocene flood history of the Tanana River, Alaska, U.S.A.: *Arctic and Alpine Research*, v. 23, no. 4, p. 392-403.

Pevé, T.L., 1951, An observation of wind-blown silt: *Journal of Geology*, v. 59, p. 399-401.

—1955, Origin of the upland silt near Fairbanks, Alaska: *Geological Society of America Bulletin*, v. 66, no. 6, p. 699-724.

—1970, Permafrost and vegetation on floodplains of subarctic rivers (Alaska), a summary, in *Ecology of the subarctic regions—Proceedings of the Helsinki symposium*, UNESCO, p. 141-142.

Pevé, T.L., and Holmes, G.W., 1964, Geology of the Mt. Hayes-D-4 Quadrangle, Alaska: U.S. Geological Survey Miscellaneous Geologic Investigations Map 1-394, scale 1:63,360, 2 sheets.

Pevé, T.L., and Reger, R.D., 1983a, Delta River area, Alaska Range, in Pevé, T.L., and Reger, R.D., eds., Guidebook to permafrost and Quaternary geology along the Richardson and Glenn Highways between Fairbanks and Anchorage, Alaska: Alaska Division of Geological & Geophysical Surveys Guidebook 1, p. 47-135.

—1983b, Middle Tanana River valley, in Pevé, T.L., and Reger, R.D., eds., Guidebook to permafrost and Quaternary geology along the Richardson and Glenn Highways between Fairbanks and Anchorage, Alaska: Alaska Division of Geological & Geophysical Surveys Guidebook 1, p. 5-45.

Ping, C.L., Boone, R.D., Clark, M.H., Packer, E.C., and Swanson, D.K., 2006, State factor control of soil formation in interior Alaska, in Chapin, F.S., III, Osowood, M.W., Van Cleve, Keith, Viereck, L.A., and Verbyla, D.L., eds., Alaska's changing boreal forest: New York, Oxford University Press, p. 21-38.

Reger, R.D., and Hubbard, T.D., 2009, Reconnaissance interpretation of 1978-1980 permafrost, Alaska Highway Corridor, Robertson River to Tetlin Junction, Alaska: Alaska Division of Geological & Geophysical Surveys Preliminary Interpretive Report 2009-6a, 4 sheets, scale 1:63,360.

Reger, R.D., Stevens, D.S.P., and Solt, D.N., 2008, Surficial geology of the Alaska Highway corridor, Big Delta and Mt. Hayes quadrangles, Alaska: Alaska Division of Geological & Geophysical Surveys Preliminary Interpretive Report 2008-3a, 48 p., 2 sheets, scale 1:63,360.

Sloan, C.E., Zenone, Chester, and Mayo, L.R., 1976, Kings along the Trans-Alaska pipeline route: U.S. Geological Survey Professional Paper 979, 31 p.

Springer, W., George, T.H., and Bell, R.M., 1976, Identification of flood hazard resulting from auflages formation in an interior Alaskan stream: Unpublished University of Alaska Geophysical Institute report for U.S. Department of Agriculture Soil Conservation Service, 12 p.

Weber, F.R., 1986, Glacial geology of the Yukon-Tanana Upland, in Hamilton, T.D., Reed, K.M., and Thoreson, R.M., eds., Glaciation in Alaska: The geologic record: Anchorage, Alaska Geological Society, p. 79-98.

Weber, F.R., and Pevé, T.L., 1961, Engineering geology problems in the Yukon-Kuskokwim lowland, Alaska, in Short Papers in the Geologic and Hydrologic Sciences 1961: U.S. Geological Survey Professional Paper 1244-D, p. 371-373